

Cerus imbridatus Haw 1821 Rev. p. 70 1821 Munha værea 20. Rev. p.66.4.15 1826 From Mexica derspriens H Ris 1829 p. 118 (pm Coultur) 1 hebs Fally De. Prot. 1828 3.471. (Hawoth) (arborespen) Egel Bost Will 1848) Exaviata De Rev: 1829 p. 118

a=tunicata Bag= inbritta p.

furiosa wend Cat Harnhey 1835

= tunicata Lehm van fordyn Oristata Salm Cast ell p.49 = inbritte firstelm Ø imbritate es the older name [Hawortt 1821 Jul Cereo O. rosen & New tab 15 has similar flowers [2" Dian) and leaves (of wary).

ody 3 lines long, fruit apparely similar to arborregrang

seems to belong to tunisate Lihon. 4. In Salar p. 250 Ju also for Synonymy Salen 1849 p.73 & 249 weber's Nant, Majors (15-12) & Sin all with Broad

Hay a Schwith

Some dain Potoni, Parry.

Brong. Marry 26h 200, 24-24 30 Carperle

Wirdlige Parry.

Wirdlige Parry.

120-24 15 Carperle Such figured of The same of the sa MISSOURI

copyright reserved

cm

BOTANICAL

GARDEN

Cyberia qualice of the Books, probelly allied to O, imbricate Op. imbrieta (Hans) &C. Prot III. 471. O. Deripians & C. Nw. p. M. (not = Orod ?)

O. Deripians & C. Nw. p. M. (nomen)

O. cristata Salum Cast well prof = melasta for of God 1849 O. rosea &C Nevery 66 t. 15 1009 ans. 1828 O convicta se ipart. Par 118 O. stelleta Sala (where) p. 48 = arbonom for Sala & Ben Bit HD.

Reinine & C. Per p. 118. 1829. O. Stationente (Los on Cet 1832 Ceres) Holand. Pfiffer O. examinate St-jant. Aw 11X O. punissa went Cit Herrol. 1835 O. autorison Egl. Wist Reg. notes 1848 DC. Port has only 111. 471: rosen x indicate de Riffer em 1835 : exavist, turi rathe submit rose att Mapelinia (meritum) Kleinie Streeipiens St AC. Rev. 1829 In Ausp: exemple de prime, Restrice. posse MISSOURI

copyright reserved cm


BOTANICAL GARDEN

realther has been seven several years gens on fair and so the constant has being and so the constant and attention and attention and attentions are attentions and attentions are attentions and attentions and attentions are attentions are attentions and attentions are attentions and attentions are attentions are attentions and attentions are attentions and attentions are attentions are attentions are attentions are attentio South St. Louis Constitution of the second of MISSOURI MOTHINGAL GARDEN MISSOURI

BOTANICAL copyright reserved cm GARDEN

Opul: i'mbrierte Han Tog Corro O. decipiens DO. ho (Synon) O cristata Selve (X var ! O, exerciata & C. Aw.) O, vosca De Aw tybes. De 10 or III, 471 ha O. rosea De O subnista De O. tosta so Rev. pl. 14 has red flows with selectory oray but smooth Ditch fit = astorems.

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved



Muntis imbrenta Can Ling Stefferry Morra elery deall 1/2 - 2/4 Chiles Cong Spins 18-6 or over 20 ! abot 15 anolum ovary (in other, 21 monly) Have alet & feel Ligh branchers voitisellet noting organion as certificant estill has degree from the planting


0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

GARDEN

MISSOURI
BOTANICAL
GARDEN

MISSOURI BOTANICAL GARDERS GEORGE ENGELMANN PAPERS




0 1 2 3 4 5 6 7 8 9 10 MIS

cm copyright reserved G A

June 1782


Opentis turirate and arbivesrens are closely allied from numerous arrounts of must conclude that lumbrienta is not taller they the northern plant, my 4-5 feet high, but the Sperimen I have seen in aultivation in Montpellier is probly 12 feel high. The flower of imbricate are generally smaller, the fruit soft, pulpy, greenish, its tubersles not very provincent, the seed larger, more irregular, with a broader and more prominent Vhaple, sweetish puly often addering to these, tuturely of the steer more prominent (justifying the name of Mistata Johich Talun give to one form) O asborsons has nother less prosecioned technolog, layer flowers, a strongly futeroled er oristete rather leather fruit, whitesom beaung dre and the regular, with a narrower, not prominent shappe. and without may pulso addening. Parry's San Luis sperime of imbritate has smaller seed with irregular, and with last prominent Roughe, approaching abording af a puly fruit. There are probably other allies speries in Mexico, mally with Town




Inteller flowers, Ach I am now unable to disting the.
The only closely ellied Mexican speries described is Op.
Kleinine with slanderer branches and much smaller flowers.

MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS


(1xe) Ker John Just John Just




He Mus. Paris April 1869 264 Gendelige, Mexito, Esper- finne, us 8. Burgen 20 Mai 1865 The stanter langer flower plant with globone fruit approaching, if not identisel esitt, my O. auboressen, seens to be O imbritate the the ATAMAN finden la on This muner Enstly soon Inspears, but den a very your joshts and on over ANts yellen 4 ovany 1/4 shih lay, with about 20 arrolae, onstite, etszeph 46" flower 2/2 shels Naul oboote spatialité pulit, 6-3 strymte ved fri Anbylobose, vry enstit Toffelat last now, before ripens) with large slightly insumers Tumble





March 27 12/6 Op. rubsitate Hægge deboundt, Efort. 1873 Thaphe always above broadest abruptly narrowed to wards the hileun and terminating in a their possely the attachement of the funitables; On lower part fradicular part of seed shaphe not as evide es above terminaling bleintly.



MISSOURI BOTANICAL GARDERS GEORGE ENGELMANN PAPERS









MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS







Openles jubirale Bourgeau, Mexist, nem City 264 his 20 may 1×65 Jud 2,5 lines
prisologiestone sim, commission, or saphe, very broad and some what por in our 2 /2 Chan Stan reed Cargo, Thick, regular Juit of 264 obvinte, smaller seeds exactly the same but seavely own 2,2 lines with







MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS



Tai manh 1869 Opuntia inbrinta Det Weber's account: common about the city of Mexiso, and generally from or nother vanegas, a little north of matchuala) the orly (and a very E good / five evond. Hank 3-6 feet Ligh, 3-4 inches that at bese. if I sperime of word brought home, is a little poor 3 inches in La meter and shows the same I hollow oglinden, with retribeted weeks as O. aubousreus, but is less compact, meshes g i Ze order, 8 mg s spirale the most prominent; mesher larger them is aubresteer, From De Webers notes, the flowers are 4 c. 4 (18 ling) Samuter, but petals 10 i number 2/2 rent. lay and 2 wide (12 l. lay gor 10 (with)
flower therefore probably supshaped _ as that of Vaporesnens. Petels obovate trumoate, erose decliralete at the truncate end, voic puple with greenisk bace Stammer previolen, 7 whitish yellowsh fruit yellen, 3 r.m i d'aunte (14 lines) globulan, Lubernlated, spinons, out spines deridous at maturity. fleshy bet not jurity, å rid. They are used av a mordant in a obvering with cuche wille In'est frist, skylly enlarget of restoring showelles toberely which are 15-20 inumb sæds vitte a Grond Asleyhte prominent rommissem

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved



GEORGE ENGELMANN PAPERS

